

**2.1.059*****Labrys potucalensis*, a bacterial strain with the capacity to degrade fluorobenzene**

**M.F. Carvalho<sup>1,\*</sup>, P. de Marco<sup>2</sup>, A.F. Duque<sup>1</sup>, C.C. Pacheco<sup>2</sup>, D.B. Janssen<sup>3</sup>, P.M. Castro<sup>1</sup>**

<sup>1</sup> CBQF/Escola Superior de Biotechnologia, Univeridade Catolica Portuguesa, Porto, Portugal

<sup>2</sup> IBMC-Instituto de Biologia Molecular e Celular, Universidade do Porto, Porto, Portugal

<sup>3</sup> Biochemical Laboratory, Groningen Biomolecular Sciences and Biotechnology Institute, University of Groningen, Groningen, Netherlands

During previous studies on the microbial degradation of fluorobenzene (FB), a pure bacterial culture with the unique capacity to utilize this compound as a sole carbon and energy source was isolated from a sediment sample collected from an industrially polluted site in northern Portugal. The isolated strain, designated F11, degrades FB via 4-fluorocatechol, with subsequent *ortho* cleavage, and also partially via catechol. It has the ability to utilize other aromatic compounds including other fluorinated compounds such as 2-fluorobenzoate, 4- fluorobenzoate and 4-fluorophenol.

Identification studies showed that strain F11 belongs to subgroup 2 of the class Alphaproteobacteria and falls within the order *Rhizobiales*. By using a polyphasic approach, which included analysis of morphological and physiological characteristics, cellular fatty acid profiling, phylogenetic analysis of the 16S rRNA gene and DNA—DNA hybridization experiments, strain F11 was identified as a novel species of the genus *Labrys*.

doi:[10.1016/j.nbt.2009.06.306](https://doi.org/10.1016/j.nbt.2009.06.306)